

References:

- Andrews B, Semler MW, Muchemwa L, et al. Effect of early resuscitation protocol on in-hospital mortality among adults with sepsis and hypotension: a randomized clinical trial. *JAMA* 2017;318(13):1233-40.
- Angus DC, Poll T. Severe sepsis and septic shock. *N Engl J Med* 2013;369:840-51.
- Annane D, Siami S, Jaber S, et al. Effects of fluid resuscitation with colloids vs crystalloids on mortality in critically ill patients presenting with hypovolemic shock: the CRISTAL randomized trial. *JAMA* 2013;310(17):1809-17.
- Bone RC, Balk RA, Cerra FB, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Chest* 1992;101(6):1644-55.
- Boyd JH, Forbes J, Nakada T, et al. Fluid resuscitation in septic shock: a positive fluid balance and elevated central venous pressure are associated with increased mortality. *Crit Care Med* 2011;39:259-265.
- Caironi P, Tognoni G, Masson S, et al. Albumin replacement in patients with severe sepsis or septic shock. *N Engl J Med* 2014;370:1412-21.
- Chowdhury AH, Cox EF, Francis ST, et al. A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and Plasma-Lyte 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 2012;256:18-24.
- Faust JS, Weingart SD. The past, present, and future of the Centers for Medicare and Medicaid Services Quality Measure SEP-1: The early management bundle for severe sepsis/septic shock. *Emerg Med Clin N Am* 2017;35:219-231.
- Genga K, Russell JA. Early liberal fluids for sepsis patients are harmful. *Crit Care Med* 2016;44:2258-62.
- Genga KR, Russell JA. How much excess fluid impairs outcome of sepsis? *Intensive Care Med* 2017;43:680-2.
- Guirgis FW, Williams DJ, Hale M, et al. The relationship of intravenous fluid chloride content to kidney function in patients with severe sepsis or septic shock. *Am J Emerg Med* 2015;33:439-43.
- Hjortrup PB, Haase N, Wetterslev J, et al. Gone fishing in a fluid trial. *Crit Care Resusc* 2016;18:55-58.
- Jaehne AK, Rivers EP. Early liberal fluid therapy for sepsis patients is not harmful: hydrophobia is unwarranted but drink responsibly. *Crit Care Med* 2016;44(12):2263-9.
- Jaffee W, Hodgins S, McGee WT. Tissue edema, fluid balance, and patient outcomes in severe sepsis: an organ systems review. *J Intensive Care Med* 2017;doi: 10.1177/0885066617742832. [Epub ahead of print].

- The Joint Commission. Specifications Manual for National Hospital Inpatient Quality Measures. Specifications Manual for National Hospital Inpatient Quality Measures v5.3. Available at: <https://www.jointcommission.org/>. Accessed January 7, 2018.
- Lee SJ, Ramar K, Park JG, et al. Increased fluid administration in the first three hours of sepsis resuscitation is associated with reduced mortality: A retrospective cohort study. *CHEST* 2014; 146:908–915 41.
- Leisman D, Wie B, Doerfler M, et al. Association of fluid resuscitation initiation within 30 minutes of severe sepsis and septic shock recognition with reduced mortality and length of stay. *Ann Emerg Med* 2016; 68:298–311 43.
- Levy MM, Dellinger RP, Townsend SR, et al. The surviving sepsis campaign: results of an international guideline-based performance improvement program targeting sepsis. *Crit Care Med* 2010;38:367-74.
- Levy MM, Rhodes A, Phillips GS, et al. Surviving sepsis campaign: association between performance metrics and outcomes in a 7.5 year study. *Crit Care Med* 2015;43:3-12.
- Maitland K, Kiguli S, Opoka RO, et al. Mortality after fluid bolus in African children with severe infection. *N Engl J Med* 2011;364(26):2483–95.
- Marik PE, Linde-Zwirble WT, Bittner EA, et al. Fluid administration in severe sepsis and septic shock, patterns and outcomes: an analysis of a large national database. *Intensive Care Med* 2017;43(5):625-32.
- Moeller C, Fleischmann C, Thomas-Rueddel D, et al. How safe is gelatin? A systematic review and meta-analysis of gelatin-containing plasma expanders vs crystalloids and albumin. *J Crit Care* 2016;35:75-83.
- Mouncey PR, Osborn TM, Power S, et al. Trial of early, goal-directed resuscitation for septic shock. *N Engl J Med* 2015;372(14):1301-11.
- Murphy CV, Schramm GE, Doherty JA, et al. The importance of fluid management in acute lung injury secondary to septic shock. *CHEST* 2009;136:102-109.
- Myburgh JA, Finfer S, Bellomo R, et al. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med* 2012;367:1901-11.
- Myburgh JA, Mythen MG. Resuscitation fluids. *N Engl J Med* 2013;369:1243-51.
- Omron EM, Omron RM. A physiochemical model of crystalloid infusion on acid-base status. *J Intensive Care Med* 2010;25(5):271-80.
- Parks Taylor S, Karvetski CH, Templin MA, et al. Initial fluid resuscitation following adjusted body weight dosing is associated with improved mortality in obese patients with suspected septic shock. *J Crit Care* 2018;43:7-12.
- Peake SL, Delaney A, Bailey M, et al. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med* 2014;371(16):1496-1506.

- Perner A, Haase N, Guttormsen AB, et al. Hydroxyethyl starch 130/0.42 versus ringer's acetate in severe sepsis. *N Engl J Med* 2012;367:124-34.
- Raghunathan K, Shaw A, Nathanson B, et al. Association between in-hospital mortality among critically ill adults with sepsis. *Crit Care Med* 2014;42:1585-1591.
- Rhodes A, Evans LE, Alhazzani W, et al. Surviving sepsis campaign: International guidelines for management of sepsis and septic shock: 2016. *Crit Care Med* 2017;45(3):486-552.
- Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 2001;345:1368-77.
- The SAFE study investigators. A comparison of albumin and saline for fluid resuscitation in the intensive care unit. *N Engl J Med* 2004;350:2247-56.
- Semler MW, Rice TW. Sepsis resuscitation: fluid choice and dose. *Clin Chest Med* 2016;37(2):241-250.
- Singer M, Deutschman CS, Seymour CW, et al. The third international consensus definitions for sepsis and septic shock (sepsis-3). *JAMA* 2016;315(8):801-10.
- Yealy DM, Kellum JA, Huang DT, et al. A randomized trial of protocol-based care for early septic shock. *N Engl J Med* 2014;370(18):1683-93.
- Young P, Bailey M, Beasley R, et al. Effect of a buffered crystalloid solution vs saline on acute kidney injury among patients in the intensive care unit: The SPLIT randomized clinical trial. *JAMA* 2015;314(16):1701-10.
- Yunos NM, Bellomo R, Hegarty C, et al. Association between a chloride-liberal vs chloride-restrictive intravenous fluid administration strategy and kidney injury in critically ill adults. *JAMA* 2012;308(15):1566-72.
- Zarychanski R, Abou-Setta AM, Turgeon AF, et al. Association of hydroxyethyl starch administration with mortality and acute kidney injury in critically ill patients requiring volume resuscitation: a systematic review and meta-analysis. *JAMA* 2013;309(7):678-88.