

References

- 1) Zingale LC, et al. "Acquired deficiency of the inhibitor of the first complement component: presentation, diagnosis, course, and conventional management." *Immunol Allergy Clin North Am*. 2006;26(4):669.
- 2) Cicardi M, et al. "Classification, diagnosis, and approach to treatment for angioedema: consensus report from the Hereditary Angioedema International Working Group." *Allergy*. 2014 May;69(5):602-16.
- 3) Bas M, et al. "Increased C-reactive protein in ACE-inhibitor-induced angioedema." *Br J Clin Pharmacol.* 2005 Feb;59(2):233-8.
- 4) Chase MP, et al. "Angioedema of the small bowel due to an angiotensin-converting enzyme inhibitor." *J Clin Gastroenterol.* 2000;31(3):254.
- 5) Vallurupalli K, et al. "MDCT features of angiotensin-converting enzyme inhibitor-induced visceral angioedema." *AJR Am J Roentgenol*. 2011;196(4):W405.
- 6) Kaplan AP, et al. "Angioedema." J Am Acad Dermatol. 2005;53(3):373.
- 7) Bas M, et al. "A randomized trial of icatibant in ACE-inhibitor-induced angioedema." *N Engl J Med*. 2015 Jan;372(5):418-25.
- 8) Straka BT, et al. "Effect of bradykinin receptor antagonism on ACE inhibitor-associated angioedema." J Allergy Clin Immunol. 2017;140(1):242.
- 9) Dean DE, et al. "Asphyxia due to angiotensin converting enzyme (ACE) inhibitor mediated angioedema of the tongue during the treatment of hypertensive heart disease." *J Forensic Sci.* 2001;46(5):1239.



INDIANA UNIVERSITY SCHOOL OF MEDICINE

Indiana University Health



References

- 10) Vleeming W, et al. "ACE inhibitor-induced angioedema. Incidence, prevention and management." Drug Saf. 1998;18(3):171.
- 11) Agah R, et al. "Angioedema: the role of ACE inhibitors and factors associated with poor clinical outcome." *Intensive Care Med.* 1997 Jul;23(7):793-6.
- 12) Brown NJ, et al. "Black Americans have an increased rate of angiotensin converting enzyme inhibitor-associated angioedema." *Clin Pharmacol Ther.* 1996;60(1):8.
- 13) Gibbs CR, et al. "Angioedema due to ACE inhibitors: increased risk in patients of African origin." Br J Clin Pharmacol. 1999 Dec;48(6):861-5.
- 14) Beltrami L, et al. "Long-term follow-up of 111 patients with angiotensin-converting enzyme inhibitor-related angioedema." *J Hypertens*. 2011;29(11):2273.
- 15) Toh S, et al. "Comparative risk for angioedema associated with the use of drugs that target the renin-angiotensin-aldosterone system." *Arch Intern Med*. 2012;172(20):1582.
- 16) Rasmussen ER, et al. "Angiotensin II receptor blockers are safe in patients with prior angioedema related to angiotensin-converting enzyme inhibitors a nationwide registry-based cohort study." *J Intern Med.* 2019;285(5):553.



